

Baryons in AdS/QCD

Deog Ki Hong,^{1,2,*} Takeo Inami,^{3,†} and Ho-Ung Yee^{4,‡}

¹*Department of Physics, Pusan National University, Busan 609-735, Korea*

²*Asia Pacific Center for Theoretical Physics,
POSTECH, Pohang 709-784, Korea*

³*Department of Physics, Chuo University, Tokyo 112-8551, Japan*

⁴*School of Physics, Korea Institute for Advanced Study, Seoul 130-012, Korea*

(Dated: February 2, 2008)

Abstract

We construct a holographic model for baryons in the context of AdS/QCD and study the spin- $\frac{1}{2}$ nucleon spectra and its couplings to mesons, taking full account of the effects from the chiral symmetry breaking. A pair of 5D spinors is introduced to represent both left and right chiralities. Our model contains two adjustable parameters, the infrared cutoff and the Yukawa coupling of bulk spinors to bulk scalars, corresponding to the order parameter of chiral symmetry. Taking the lowest-lying nucleon mass as an input, we calculate the mass spectrum of excited nucleons and the nucleon couplings to pions. The excited nucleons show a parity-doubling pattern with smaller pion-nucleon couplings.

PACS numbers: 11.25.Tq, 12.38.Bx, 11.25.Wx, 11.10.Kk

*E-mail: dkhong@pusan.ac.kr

†E-mail: inami@phys.chuo-u.ac.jp

‡E-mail: ho-ung.yee@kias.re.kr

I. INTRODUCTION

Quantum chromodynamics (QCD) is now firmly believed to be the theory of strong interactions. It describes the strongly interacting particles at high energies or in deep-inelastic scattering extremely well, where the weak coupling perturbation is reliable. QCD at low energies is on the other hand strongly coupled and highly nonlinear. So far it has precluded any analytic solutions. The test of QCD at low energies, therefore, has relied on either indirect evidences, based on chiral perturbation theory, or on lattice calculations for limited cases. Any reliable approximate methods to solve QCD deserve a careful study, since we may gain physical insight from them into the low energy dynamics of QCD.

Recent discovery of the correspondence between a string theory in anti-de Sitter (AdS) space and the conformal field theory (CFT) in its boundary has established a new paradigm for understanding strongly coupled gauge theories [1, 2, 3]. The holographic dual of strongly coupled gauge theory corresponds to the low energy limit or the supergravity limit of string theory in the large 't Hooft coupling limit, which therefore provides a solvable scheme for QCD. In this scheme, called AdS/QCD, taking the number of color very large, one approximates QCD to be conformally invariant and thus its coupling and the anomalous dimensions of composite operators are taken to remain constant for all scales higher than the QCD scale, Λ_{QCD} , which is then introduced by hand. AdS/QCD will break down at high energies, where QCD is asymptotically free. However, it was shown to give reasonable results for the meson sector of low energy QCD [4, 5, 6, 7, 8, 9, 10]. This scheme has also been successfully applied for technicolor theories [11, 12, 13, 14, 15], which are otherwise very hard to solve.

As baryons are essential ingredients in QCD, it is important to extend the above models for mesons to include baryons. In this paper we construct a model for low-lying spin- $\frac{1}{2}$ baryons in the context of AdS/QCD and study their spectra and couplings to mesons, taking full account of the effects from chiral symmetry breaking. Some efforts have been already made to study the baryon spectrum of high spin/mass resonances [16]. However, the effects of chiral symmetry breaking in QCD were not exploited. Since such effects are important for low spin/mass baryons, our analysis is a first step to resolve the discrepancy. We also point out that the recent model of Sakai and Sugimoto from string theory [6] contains baryons as solitonic objects, though its detailed study has not been done yet. We discuss the relation

between our approach and theirs in section V.

II. HOLOGRAPHIC BARYONS AND CHIRAL SYMMETRY

The low-lying states of baryons form an even-parity octet or decuplet under the flavor $SU(3)$ symmetry for three light u, d, s quarks. The excited states of baryons form singlets, octets or decuplets but nearly degenerate in parity. One can in principle calculate in QCD the spectra of ground states and radially excited states of the baryon octet and decuplet from the correlation functions of following composite operators,

$$B_j^i = \frac{1}{\sqrt{3}} \left(q^i q^k q^l \epsilon_{jkl} - \frac{1}{3} \delta_j^i q^m q^k q^l \epsilon_{mkl} \right) \quad \text{and} \quad \Delta^{ijk} = \frac{1}{\sqrt{6}} (q^i q^j q^k + \text{cyclic perm.}), \quad (1)$$

which have the same quantum number of the baryon octet and decuplet, respectively. (The orbitally excited states will contain positive powers of derivatives.) In Eq. (1) the italic indices denote the quark flavors u, d, s and we have suppressed both Dirac indices and color indices of quarks, keeping the color indices completely anti-symmetric to form a color singlet.

According to the AdS/CFT correspondence, the spin- $\frac{1}{2}$ baryon operator at the boundary will correspond to a bulk Dirac field [17], while the spin- $\frac{3}{2}$ baryon operator to a bulk Rarita-Schwinger field [18]. Note that spinors in 5D bulk are always Dirac spinors. In this paper we consider the spin- $\frac{1}{2}$ baryons only for simplicity. It is straightforward to extend our analysis to spin- $\frac{3}{2}$ baryons. The key idea in our model is to introduce a pair of bulk spinors to describe the boundary Dirac spinor. This predicts the parity-doublet pattern of excited baryons, explaining the observed parity-doubling in excited baryons from the holographic picture even though the ground state still breaks the chiral symmetry.¹

The ground state spin- $\frac{1}{2}$ baryons will be massless if the chiral symmetry of QCD is unbroken.² As they are composed of chiral quarks, their left-handed and right-handed components separately fall into $(8, 1)$ and $(1, 8)$ or $(3, \bar{3})$ and $(\bar{3}, 3)$ under the chiral symmetry $SU(3)_L \times SU(3)_R$. Another way of describing this is to introduce a pair of spin- $\frac{1}{2}$ chiral baryon operators $\mathcal{O}_L$ and $\mathcal{O}_R$ which transform as $(8, 1)$ and $(1, 8)$ or $(3, \bar{3})$ and $(\bar{3}, 3)$ under

¹ See [19] for a similar explanation on parity-doubling.

² For two flavors the massless spin- $\frac{1}{2}$ baryons are fundamental under the flavor symmetry $SU(2)_L \times SU(2)_R$ and saturate the axial anomaly as originally shown by 't Hooft [20]. However for three flavors they do not saturate the anomalies of the microscopic theory. One therefore suspects the chiral symmetry has to be broken for three flavors.

chiral symmetry. A natural mechanism for nucleon masses would be then the mass coupling between them by both explicit and spontaneous chiral symmetry breaking.

The above picture for the effects of chiral symmetry breaking on the baryon spectra works well for the up and down quarks but not for the strange quark due to the large strange quark mass. In this paper we therefore focus on the two-flavor case with vanishing current quark mass, where the effect from the quark-antiquark condensate dominates.

To account for the chiral symmetry and its breaking in holographic QCD, we adopt the model in [7, 8] and take a slice of the AdS metric as

$$ds^2 = \frac{1}{z^2} (-dz^2 + \eta^{\mu\nu} dx_\mu dx_\nu), \quad \epsilon \leq z \leq z_m, \quad (2)$$

where $z_m = 1/\Lambda_{\text{QCD}}$ is the IR boundary and $z = \epsilon \ll 1$ is the UV boundary, and $\eta^{\mu\nu} = \text{diag}(+ - - -)$ is the four-dimensional Minkowski metric. We introduce a bulk scalar field X_i^j , bi-fundamental under $SU(2)_L \times SU(2)_R$, which corresponds to the composite operator $\bar{q}_{iL} q_R^j$ at the boundary, described by the action

$$S_X = \int d^5x \sqrt{g} \text{Tr} \left[|DX|^2 - M_5^2 |X|^2 - \frac{1}{2g_5^2} (F_L^2 + F_R^2) \right] \quad (3)$$

where M_5 is the mass of the bulk scalar field X and the covariant derivative $D_M X = \partial_M X - iA_{LM} X + iX A_{RM}$. The bulk mass is determined by the relation [2, 3], $\Delta_0 (\Delta_0 - 4) = M_5^2$, where Δ_0 is the scaling dimension of the boundary operator $\bar{q}_{iL} q_R^j$. (For QCD $\Delta_0 = 3$.) F_L and F_R are the field strength tensors of the $SU(2)_L \times SU(2)_R$ bulk gauge fields A_L and A_R , respectively. We take the five dimensional coupling $g_5 = 2\pi$ to match the two-point correlation functions of vector currents in QCD at the ultra-violet (UV) region [7, 8].

When evaluated for the classical solutions to the bulk equations of motion, the vector and axial vector gauge fields, defined as $V = (A_L + A_R)/\sqrt{2}$ and $A = (A_L - A_R)/\sqrt{2}$, at the UV boundary couple to vector and axial vector currents $J_V^{a\mu} = \bar{q}\gamma^\mu t^a q$ and $J_A^{a\mu} = \bar{q}\gamma^\mu \gamma_5 t^a q$ of quarks, respectively. (The $SU(2)$ generators $t^a = \frac{1}{2}\sigma^a$, where σ^a 's are Pauli matrices.) The classical solution of the bulk field X is given as

$$X_0(z) = \frac{1}{2} M z + \frac{1}{2} \sigma z^3, \quad (4)$$

where the constants M and $\sigma = \langle \bar{q}_L q_R \rangle$ correspond to the bare mass and the chiral condensate of QCD, respectively. In the holographic dual of QCD the axial gauge fields therefore get a 5D mass

$$m_A^2 = g_5^2 \frac{|X_0|^2}{z^2}, \quad (5)$$

where we take $M \rightarrow 0$ in the chiral limit. The spectra and interactions from this theory have successfully reproduced the observed low energy meson physics [4, 5, 6, 7, 8, 9, 10].

Since the spin- $\frac{1}{2}$ chiral baryon operators $\mathcal{O}_L$ and $\mathcal{O}_R$ have different representations under $SU(2)_L \times SU(2)_R$, it is not possible to assign a single 5D spinor in the bulk for both of them. We must instead introduce a pair of 5D spinors N_1 and N_2 , corresponding to $\mathcal{O}_L$ and $\mathcal{O}_R$, respectively, which transform as $(2, 1)$ and $(1, 2)$ under $SU(2)_L \times SU(2)_R$.

The bulk action for the Dirac spinor N_1 (and similarly for N_2) is then given as

$$S_{N_1} = \int d^5x \sqrt{g} \left\{ \frac{i}{2} \bar{N}_1 e_A^M \Gamma^A \nabla_M N_1 - \frac{i}{2} \left(\nabla_M^\dagger \bar{N}_1 \right) e_A^M \Gamma^A N_1 - m_5 \bar{N}_1 N_1 \right\} \quad (6)$$

where m_5 is the mass of the bulk spinor, e_M^A is the vielbein, satisfying $g_{MN} = e_M^A e_N^B \eta_{AB}$, and the Dirac matrices Γ^A 's ($A = 0, 1, 2, 3, 5$) satisfy $\{\Gamma^A, \Gamma^B\} = 2\eta^{AB}$. Fixing the gauge for the Lorentz transformation, we take the vielbeins

$$e_M^A = \frac{1}{z} \eta_M^A. \quad (7)$$

In terms of the spin connection ω_M^{AB} and the gauge fields we have introduced the Lorentz and gauge covariant derivative

$$\nabla_M = \partial_M + \frac{i}{4} \omega_M^{AB} \Gamma_{AB} - i(A_L^a)_M t^a, \quad (8)$$

where the Lorentz generator $\Gamma^{AB} = \frac{1}{2i}[\Gamma^A, \Gamma^B]$ for spinor fields. The non-vanishing components of the spin connection are

$$\omega_M^{5A} = -\omega_M^{A5} = \frac{1}{z} \delta_M^A. \quad (9)$$

By comparing the correlators for a free theory, or by looking at the UV asymptotic of wave-functions, one finds the mass of the $(d+1)$ -dimensional bulk Dirac spinor, corresponding to a boundary operator of scaling dimension Δ ,

$$m_5^2 = \left(\Delta - \frac{d}{2} \right)^2. \quad (10)$$

The sign of m_5 will be related to the chirality of the boundary operator [17, 21]. Extremizing the action (6) with respect to $\bar{N}_1$, we get

$$(ie_A^M \Gamma^A \nabla_M - m_5) N_1 = 0 \quad \text{and} \quad [\delta \bar{N}_1 e_A^5 \Gamma^A N_1]_\epsilon^{z_m} = 0. \quad (11)$$

Since the classical solution to the bulk equation of motion will be a source to the corresponding boundary operator in the AdS/CFT correspondence, when evaluated at the UV brane, it is convenient to decompose the bulk spinor as, with the 4D chirality projectors,

$$N_1 = N_{1L} + N_{1R}, \quad (12)$$

where $i\Gamma^5 N_{1L} = N_{1L}$ and $i\Gamma^5 N_{1R} = -N_{1R}$. (The four dimensional $\gamma^5 = i\Gamma^5$.) Then, the boundary terms in (11) become

$$\left[z\delta\bar{N}_{1L}N_{1R} - z\delta\bar{N}_{1R}N_{1L} \right]_\epsilon^{z_m} = 0. \quad (13)$$

We now Fourier-transform the bulk spinor as

$$f_{1L,R}(p, z) \psi_{1L,R}(p) = \int d^4x N_{1L,R}(x, z) e^{ip \cdot x}, \quad (14)$$

where the 4D spinors satisfy with $\psi_{1L} = \gamma^5 \psi_{1L}$ and $\psi_{1R} = -\gamma^5 \psi_{1R}$

$$\not{p} \psi_{1L,R}(p) = |p| \psi_{1R,L}(p) \quad (15)$$

and $f_{1L,R}$ satisfy

$$\left(\partial_z - \frac{2+m_5}{z} \right) f_{1L} = -|p| f_{1R} \quad (16)$$

$$\left(\partial_z - \frac{2-m_5}{z} \right) f_{1R} = |p| f_{1L}. \quad (17)$$

Normalizable solutions with eigenvalues $|p|$ and appropriate boundary conditions that we will specify shortly give us 4D propagating (Dirac) spinors ψ_1 with mass $|p|$. For $|p| = 0$, we may obtain a chiral fermion. Eliminating f_{1R} in (16) with Eq. (17), we get

$$\left(\partial_z^2 - \frac{4}{z} \partial_z + \frac{6+m_5-m_5^2}{z^2} \right) f_{1L} = -p^2 f_{1L}. \quad (18)$$

Similarly,

$$\left(\partial_z^2 - \frac{4}{z} \partial_z + \frac{6-m_5-m_5^2}{z^2} \right) f_{1R} = -p^2 f_{1R}. \quad (19)$$

Near $z = \epsilon$ the solution behaves as

$$f_{1L} \simeq c_1(1+2m_5)z^{2+m_5} + c_2|p|z^{3-m_5}, \quad f_{1R} \simeq c_1|p|z^{3+m_5} - c_2(1-2m_5)z^{2-m_5}. \quad (20)$$

Note that each component of f_{1L} comes along with a corresponding component in f_{1R} .

In our case the scaling dimension of the baryon composite operator is $\Delta = 9/2$ and thus $m_5 = \pm 5/2$. In 5D the spinor mass term is real but has a sign ambiguity. If we take $m_5 > 0$, z^{2+m_5} is always normalizable, but z^{2-m_5} is not normalizable. On the other hand, if we take $m_5 < 0$, z^{2-m_5} is normalizable and z^{2+m_5} is not normalizable. In this paper we choose the sign of m_5 such that the left-handed chiral zero mode from N_1 , before introducing chiral symmetry breaking, corresponds to the massless left-handed nucleons. Setting $|p| = 0$ in (20) and requiring a normalizable left-handed mode, we see that $m_5 = \frac{5}{2}$ must be positive for N_1 . We then have to take $m_5 = -\frac{5}{2}$ for N_2 to get a right-handed zero mode from N_2 .

Another way of arriving at the same conclusion is the following. N_1 is a holographic description of the left-handed chiral operator $\mathcal{O}_L$. For $m_5 > 0$, the dominant component as we approach $z = \epsilon \ll 1$ is $f_{1R} \sim z^{2-m_5}$ and according to AdS/CFT dictionary, this acts as a source for $\mathcal{O}_L$. Note that chirality of f_1 is kinematically selected at the boundary. Since f_1 and $\mathcal{O}_L$ have the same representation under the chiral symmetry, the coupling must be $\bar{f}_1 \mathcal{O}_L + \text{h.c.}$, which does not vanish only if $f_1 = f_{1R}$. Had m_5 been negative, we would have f_{1L} instead of f_{1R} near the boundary, which would give wrong chirality to $\mathcal{O}_L$.

To allow a left-handed zero mode for N_1 , we impose the following boundary conditions, consistent with Eq. (13),

$$\lim_{\epsilon \rightarrow 0} N_{1L}(\epsilon) = 0 \text{ (normalizability)} \quad \text{and} \quad N_{1R}(z_m) = 0. \quad (21)$$

The remaining boundary conditions for $N_{1L}(z_m)$ and $N_{1R}(\epsilon)$ are then determined by the equations of motion in (11), which become

$$\left(\partial_z - \frac{2+m_5}{z} \right) N_{1L} + i \not{\partial} N_{1R} = 0 \quad \text{and} \quad \left(\partial_z - \frac{2-m_5}{z} \right) N_{1R} - i \not{\partial} N_{1L} = 0, \quad (22)$$

where $\not{\partial} = \gamma^\mu \partial_\mu$ and γ^μ 's are the 4D gamma matrices. The boundary conditions for N_2 are similar except for interchanging $L \leftrightarrow R$ and $m_5 \leftrightarrow -m_5$.

The normalizable zero mode ($p^2 = 0$) for a free spinor with $m_5 > 1/2$ is $f_{1L} = c_1 z^{2+m_5}$ and the nonzero modes ($p^2 \neq 0$) satisfy a Bessel equation and $f_{1L,R}(p, z) = c'_1 z^{\frac{5}{2}} J_{m_5 \mp \frac{1}{2}}(|p|z)$, where $J_n(x)$ is the Bessel function of the first kind. The spectra of the excited Kaluza-Klein tower are given as the zeros of the Bessel function, $J_{m_5 + \frac{1}{2}}(|p|z_m) = 0$, as we impose $N_{1R}(z_m) = 0$. (The spectrum of non-interacting fermionic operators has been considered in [16, 21], and for the string theory context in [22, 23].)

Similarly, we introduce another bulk spinor N_2 of mass $-m_5 = -\frac{5}{2}$ for the right-handed components of the boundary baryon operator. If we take the normalizable modes of $N_{2R} \equiv$

$\frac{1}{2}(1 - i\Gamma^5)N_2$ not to vanish at the IR brane but $N_{2L}(z_m) = 0$, then they will have exactly same spectra as those of the normalizable N_1 , except the chirality of the zero mode.

The 4D parity exchanges $\mathcal{O}_L \leftrightarrow \mathcal{O}_R$. Hence its natural holographic extension in 5D would be flipping N_1 into N_2 and vice versa, together with the usual 4D parity operation. It is also consistent with the fact that 5D spinor mass term flips its sign under the parity. The even-parity massive baryons in the chiral symmetry limit will therefore correspond to $(N_1 + N_2)/\sqrt{2}$, while the odd-parity baryons to $(N_1 - N_2)/\sqrt{2}$.

Since the bulk scalar flips the 4D chirality, it naturally introduces a gauge-invariant Yukawa coupling between two bulk spinors,

$$\mathcal{L}_{\text{int}} \ni -g (\bar{N}_2 X N_1 + \text{h.c.}) , \quad (23)$$

which becomes in the chiral limit $M \rightarrow 0$

$$- \frac{1}{2} (g \sigma) z^3 \bar{N}_2 N_1 + \text{h.c.} \quad (24)$$

The zero modes of N_1 and N_2 become massive through the Yukawa interaction (23) when the chiral symmetry is broken or $\sigma \neq 0$. Otherwise they remain massless in the chiral symmetry limit, consistent with the 't Hooft anomaly matching condition. They will form a nucleon doublet under the isospin $\text{SU}(2)_I \subset \text{SU}(2)_L \times \text{SU}(2)_R$. The Yukawa interaction also mixes two towers of massive modes from each of N_1 and N_2 , and breaks their degeneracy into the parity doublet pattern.

III. NUMERICAL ANALYSIS

We have constructed a holographic QCD model for spin- $\frac{1}{2}$ baryons. The model contains two parameters, z_m (the IR cutoff scale) and $g\sigma$. Note that σ enters independently in the meson sector. We identify the lowest mass eigenstate as the nucleon isospin doublet (p, n) . We take its mass 0.94 GeV as an input to fix g , given z_m and σ . We first use the parameters of previous analysis of the meson sector [7, 8], $z_m = (0.33 \text{ GeV})^{-1}$ and $\sigma = \frac{\sqrt{2}\xi}{g_5 z_m^3}$ with $3.4 \leq \xi \leq 4.0$ to calculate the spectrum of excited resonances.

We also treat z_m as a parameter of our model and try to fit z_m and g from the first resonance, Roper N(1440) of mass 1.44 GeV, as well as the nucleons. Our best fit is for $z_m = (0.205 \text{ GeV})^{-1}$ and $g = 14.4$. In this case, baryons are assumed to have a different IR cutoff scale than mesons.

Another important possibility that we don't pursue in this paper is to relax $\Delta = \frac{9}{2}$ for the dimension of spin- $\frac{1}{2}$ baryon operators. We would certainly expect non-vanishing anomalous dimension in strongly coupled QCD. Because we are not aware of any reliable method to calculate it in our scheme, it would be interesting to take it as an additional fitting parameter.

As N_1 and N_2 are coupled with each other, the Kaluza-Klein (K-K) mode equations become

$$\begin{pmatrix} \partial_z - \frac{\Delta}{z} & -\frac{1}{2}g\sigma z^2 \\ -\frac{1}{2}g\sigma z^2 & \partial_z - \frac{4-\Delta}{z} \end{pmatrix} \begin{pmatrix} f_{1L} \\ f_{2L} \end{pmatrix} = -|p| \begin{pmatrix} f_{1R} \\ f_{2R} \end{pmatrix},$$

$$\begin{pmatrix} \partial_z - \frac{4-\Delta}{z} & \frac{1}{2}g\sigma z^2 \\ \frac{1}{2}g\sigma z^2 & \partial_z - \frac{\Delta}{z} \end{pmatrix} \begin{pmatrix} f_{1R} \\ f_{2R} \end{pmatrix} = |p| \begin{pmatrix} f_{1L} \\ f_{2L} \end{pmatrix}, \quad (25)$$

with the boundary conditions $f_{1R}(z_m) = f_{2L}(z_m) = 0$. Eigenfunctions should be normalizable with respect to the measure $\frac{dz}{z^4}$ over $z \in [0, z_m]$.

TABLE I is the result of our numeric analysis using the shooting method. If we adopt the value of $z_m = (0.33 \text{ GeV})^{-1}$ from the meson sector, we do not seem to reproduce observed spin- $\frac{1}{2}$ baryon spectrum well. This may indicate baryons have a different IR cutoff than mesons, or we should take account of possible anomalous dimension to $\Delta = \frac{9}{2}$.

$z_m(\text{GeV}^{-1})$	g	(p,n)(GeV)	N(1440)	N(1535)	3rd	4th	5th	6th
$(0.33)^{-1*}$	8.67	0.94*	2.14	2.24	3.25	3.30	4.35	4.36
$(0.205)^{-1}$	14.4	0.94*	1.44*	1.50	2.08	2.12	2.72	2.75

TABLE I: Numerical result for spin- $\frac{1}{2}$ baryon spectrum. * indicates an input and we used $\sigma = \frac{\sqrt{2}\xi}{g_5 z_m^3}$ with $3.4 \leq \xi \leq 4$.

In the case of tunable z_m , we find $z_m = (0.205 \text{ GeV})^{-1}$ and $g = 14.4$ is the best fit, nicely fitting 1.44 GeV for N(1440) and predicting 1.50 GeV for N(1535) quite well. Note that this is also consistent with the parity of N(1440) ($P = +1$) and N(1535) ($P = -1$) since our holographic picture predicts a parity-doublet pattern. After N(1535), the next parity-doublet resonance is (2.08, 2.12) GeV, which is quite off from the data. We suspect our assuming zero anomalous dimension might be responsible for the discrepancy.

In higher resonances, the mass splitting in parity doublets will decrease as the mixing due to the chiral symmetry breaking would be negligible, compared to the large original

K-K masses. It is also worthwhile to comment that the fitted values of g are in the region of validity of our model. The effective mass from Table I has the maximum strength $\frac{1}{2}g\sigma z_m^3$, whose value is less than or comparable to m_5 .

IV. NUCLEON-PION COUPLING

The low energy dynamics of hadrons is well described by chiral Lagrangian, where the baryon interaction with pions is dictated by the chiral symmetry. If we expand the chiral Lagrangian in powers of pions, the interaction for the nucleon doublet $N = (p, n)^T$ becomes

$$\mathcal{L}_{\pi NN} = \frac{g_A}{F_\pi} \bar{N} \gamma^\mu \gamma_5 t^a N \cdot \partial_\mu \pi^a + \dots, \quad (26)$$

where g_A and F_π are the nucleon axial coupling and the pion decay constant, respectively. Using the equation of motion, the leading term in the interaction Lagrangian (26) can be rewritten as

$$-ig_{\pi NN} \bar{N} \gamma_5 \sigma^a N \pi^a \quad (27)$$

with $g_{\pi NN} = g_A m_N / F_\pi$, the Goldberger-Treiman relation. The nucleon-pion coupling is measured accurately, $g_{\pi NN} \simeq 13$, from the low energy pion-nucleon scattering.

In our model of holographic QCD we have introduced bulk gauge fields $A_{L,R}$ together with bulk scalars X and bulk spinors $N_{1,2}$. Among the solutions to the equations of motion of those fields the normalizable modes will be mapped to the physical states in QCD by the AdS/CFT correspondence. To identify the 4D fields correctly we need to diagonalize the 4D kinetic terms after decomposing the bulk fields with the eigen modes of the 5D equations of motion.

Expanding the 5D Lagrangian for the bulk gauge fields up to quadratic terms, we have

$$\mathcal{L}_G \simeq -\frac{1}{2g_5^2 z} (\partial_\mu V_\nu - \partial_\nu V_\mu)^2 - \frac{1}{2g_5^2 z} (\partial_\mu V_5)^2 - \frac{1}{g_5^2 z} \partial_z V_\mu \partial^\mu V_5 + (V_M \leftrightarrow A_M). \quad (28)$$

We see that there are cross terms in the 4D kinetic Lagrangian which mix between 4D components of (axial) vector mesons and their 5th components. The 5th components of 5D bulk gauge fields provide the longitudinal components of the 4D massive vector mesons. The axial vector bosons mix further with the phase of the bi-fundamental scalar X , which can be seen easily, if we expand its Lagrangian, taking $X(x, z) = v(z)e^{iP}$,

$$\mathcal{L}_X = \text{Tr}(|DX|^2 - M_5^2 |X|^2) = \frac{v^2}{z^2} \text{Tr} \left(\partial_\mu P - \sqrt{2} A_\mu \right)^2 - \frac{v^2}{z^2} \text{Tr} \left(\partial_z P - \sqrt{2} A_5 \right)^2 + \dots. \quad (29)$$

Now we introduce gauge-fixing terms, which will remove the cross terms,

$$\mathcal{L}_{G.F.} = -\frac{z^4}{2\xi_V} \left[\partial^\mu V_\mu - \xi_V z \partial_z \left(\frac{V_5}{z} \right) \right]^2 - \frac{z^4}{2\xi_A} \left[\partial^\mu A_\mu - \xi_A z \partial_z \left(\frac{A_5}{z} \right) + 2\sqrt{2}g_5^2 \frac{\xi_A}{z^2} v^2 P \right]^2 \quad (30)$$

In the unitary gauge ($\xi_{V,A} \rightarrow \infty$) V_5 and the linear combination $z\partial_z(A_5/z) - 2\sqrt{2}v^2g_5^2P/z^2$ become infinitely massive and decouple from the theory. However, we note that its orthogonal combination of A_5 and P remain massless, if

$$z\partial_z \left(\frac{A_5}{z} \right) - 2\sqrt{2} \frac{v^2g_5^2}{z^2} P = 0. \quad (31)$$

The kinetic terms for the bulk gauge and scalar fields become in the unitary gauge

$$\begin{aligned} \mathcal{L}_K = & -\frac{1}{2g_5^2} V_\mu^a (-g^{\mu\nu} \partial^2 + \partial^\mu \partial^\nu) V_\nu^a - \frac{1}{2g_5^2} A_\mu^a \left[-g^{\mu\nu} \left(\partial^2 - \frac{g_5^2 v^2}{z^2} \right) + \partial^\mu \partial^\nu \right] A_\nu^a \\ & + \frac{z^4}{2g_5^2} \partial_\mu A_5^a \partial^\mu A_5^a + \frac{1}{2} v^2 z^2 \partial_\mu P^a \partial^\mu P^a - \frac{1}{2} v^2 z^2 \left(\partial_z P^a - \sqrt{2} A_5^a \right)^2. \end{aligned} \quad (32)$$

With the condition (31) the kinetic terms of A_5 and P can be rewritten as

$$\mathcal{L}_K(A_5, P) = \frac{z^4}{2g_5^2} (\partial_\mu A_5)^2 + \frac{z^8}{8g_5^2 v^2} \left[\partial_z \left(\frac{\partial_\mu A_5}{z} \right) \right]^2 - \frac{v^2 z^2}{8g_5^4} \left[\partial_z \left(\frac{z^3}{v^2} \partial_z \left(\frac{A_5}{z} \right) \right) - 4g_5^2 A_5 \right]^2 \quad (33)$$

Since the massless combination of A_5 and P should map to pions of QCD, the Nambu-Goldstone bosons associated with the spontaneous chiral symmetry breaking, we take an Ansatz

$$A_5(x, z) = f_0(z) \pi(x), \quad (34)$$

where $\pi(x)$ is identified as the pion field. Then, the last term in (33) should vanish and

$$\partial_z \left[\frac{z^3}{v^2} \partial_z \left(\frac{f_0}{z} \right) \right] - 4g_5^2 f_0 = 0. \quad (35)$$

In order to have a canonical kinetic term for pions we take the normalization of $f_0(z)$

$$1 = \int_0^{z_m} dz \left[\frac{1}{2g_5^2 z} f_0^2 + \frac{z^3}{8v^2 g_5^4} \left(\partial_z \left(\frac{f_0}{z} \right) \right)^2 \right] \quad (36)$$

Once the pion fields are correctly identified, it is straightforward to find their couplings to nucleons and their excited states. In holographic QCD pions couple to nucleons through the (bulk) gauge coupling and the Yukawa coupling Eq. (23),

$$\mathcal{L}_{\text{int}}(\mathcal{N}, A_5) = -i \frac{z}{\sqrt{2}} \bar{\mathcal{N}} \gamma^5 A_5 \tau_3 \mathcal{N} - \frac{gz^3}{2\sqrt{2}v} \bar{\mathcal{N}} \gamma^5 \partial_z \left(\frac{A_5}{z} \right) \tau_2 \mathcal{N}, \quad (37)$$

where we used the relation (31) and the Pauli matrices $\tau_{2,3}$ act on the doublet of 5D spinors, $\mathcal{N} = (N_1, N_2)^T$. The 5D spinors are decomposed in terms of the K-K modes of (25) as

$$\mathcal{N}_L = \sum_n \begin{pmatrix} f_{1L}^{(n)}(z)\psi_L^{(n)}(x) \\ f_{2L}^{(n)}(z)\psi_L^{(n)}(x) \end{pmatrix}, \quad \mathcal{N}_R = \sum_n \begin{pmatrix} f_{1R}^{(n)}(z)\psi_R^{(n)}(x) \\ f_{2R}^{(n)}(z)\psi_R^{(n)}(x) \end{pmatrix} \quad (38)$$

with the normalization for each modes

$$\int_0^{z_m} \frac{dz}{z^4} \left(|f_{1L}^{(n)}|^2 + |f_{2L}^{(n)}|^2 \right) = 1 = \int_0^{z_m} \frac{dz}{z^4} \left(|f_{1R}^{(n)}|^2 + |f_{2R}^{(n)}|^2 \right). \quad (39)$$

We then find the pion-nucleon-nucleon coupling for the lowest-lying nucleons

$$g_{\pi NN} = \int_0^{z_m} \frac{dz}{2\sqrt{2}z^4} \left[f_0 \left(f_{1L}^{(1)*} f_{1R}^{(1)} - f_{2L}^{(1)*} f_{2R}^{(1)} \right) - \frac{gz^2}{2vg_5^2} \partial_z \left(\frac{f_0}{z} \right) \left(f_{1L}^{(1)*} f_{2R}^{(1)} - f_{2L}^{(1)*} f_{1R}^{(1)} \right) \right] \quad (40)$$

Similarly one finds the couplings for the pion-nucleon-Roper coupling and other couplings with excited states. Numerical results for the couplings are presented in Table II.

For parity-even states, the eigenfunctions have the properties $f_{1L}^{(n)} = f_{2R}^{(n)}$ and $f_{1R}^{(n)} = -f_{2L}^{(n)}$, while parity-odd states satisfy $f_{1L}^{(n)} = -f_{2R}^{(n)}$ and $f_{1R}^{(n)} = f_{2L}^{(n)}$. The lowest-lying ($n = 1$) nucleons, (p, n), are parity even, and the excited resonances appear as parity-doublet (N^+, N^-) with the parity-even states (N^+) slightly lighter than the parity-odd states (N_-), as shown in Table I. These are all verified numerically.

Similarly to the pion coupling to the lowest-lying nucleons, Eq. (27), the coupling of the pion to the nucleon N and the first-excited resonance $N(1440)$ is given as $ig_{\pi NN_{1440}} \bar{N} \sigma^a \gamma^5 N_{1440} \pi^a + \text{h.c.}$, whereas the coupling to the parity-odd excitation $N(1535)$ as $ig_{\pi NN_{1535}} \bar{N} \sigma^a N_{1530} \pi^a + \text{h.c.}$. Our K-K reduction reproduces precisely the above interactions that are consistent with parity.

As we vary z_m away from the fit from the meson sector, the value of the chiral condensate σ in (4) must be determined by other means. We adopt the fit in [8], $\sigma = \frac{\sqrt{2}\xi}{g_5 z_m^3}$ with $3.4 \leq \xi \leq 4$ for all z_m . TABLE II is our prediction for these couplings when $z_m = (0.205 \text{ GeV})^{-1}$. For $z_m = (0.33 \text{ GeV})^{-1}$ we find $g_{\pi NN} = 4.35, 3.78, 3.30$ for $\xi = 3.4, 3.7, 4.0$, respectively.

V. RELATION TO BARYONS AS SKYRMIONS

In AdS/QCD we have gauge fields living in the bulk AdS whose dynamics is dual to the meson sector of QCD, including pions and higher resonances. As is well-known, the pion

$z_m(\text{GeV}^{-1})$	ξ	$g_{\pi NN}$	$g_{\pi NN_{1440}}$	$g_{\pi NN_{1535}}$
$(0.205)^{-1}$	3.4	7.34	-0.025	2.04
$(0.205)^{-1}$	3.7	6.52	0.30	1.88
$(0.205)^{-1}$	4.0	5.84	0.56	1.74

TABLE II: Numerical results for pion-nucleon-nucleon couplings.

sector can have a topological skyrmion solution, which is interpreted as a baryon, and its fermionic character comes from a topological WZW term upon quantization.

The QCD skyrmion appears as an “instanton particle” in the bulk AdS [6, 24]. Neglecting the time direction, AdS₅ has 4 spatial directions (R^3+z -direction) and we can have a gauge field configuration along these 4 directions which simply looks like an ordinary instanton configuration. It is a static, particle-like, Minkowski object in AdS₅ spacetime. Its fermionic nature is introduced by a 5d Chern-Simons term in AdS₅, which is again dual to the WZW term in QCD. These terms should exist to match chiral anomalies. If we identify Nambu-Goldstone pions as Wilson lines of the broken axial gauge fields, the topological instanton charge in AdS₅ is identical to the winding number of skyrmions in QCD [24, 25], confirming the correspondence.

In QCD, we may describe baryons as either skyrmions of the chiral Lagrangian or we may introduce separate baryon fields independently and couple them to mesons. The latter seems more effective and convenient. It is easier to identify and realize chiral symmetry property in the latter approach.

We can take the same effective approach in the holographic dual of QCD. We describe “instanton particles” with a 5d Chern-Simons term in AdS₅ by effective local spinor fields, and couple them to gauge fields representing the meson sector of QCD. Aspects of chiral symmetry and its breaking are more manifest in this effective theory model as we have studied in this work.

Another justification for this approach is the following. As the axial gauge symmetry is broken in AdS₅ spacetime, its instantons representing holographic baryons tend to shrink to zero size, eventually go beyond the validity of our effective theory. As they become point-like small instantons, we may introduce an effective baryon field to describe them. In a more detailed model like $D4-D8-\overline{D8}$ system [6], the size of the instantons is stabilized,

and the effective field description would be reliable in low energy limit. It also explains the appearance of two 5D spinors in our model; as we have $SU(2)_L \times SU(2)_R$ gauge fields, N_1 describes $SU(2)_L$ “instanton particles” and N_2 corresponds to $SU(2)_R$ “instanton particles”. Their charges arise from quantization of collective coordinates.

VI. CONCLUSION AND OUTLOOK

We have constructed a holographic model of QCD to study the spectrum of nucleons and their coupling to pions. In our approach it is necessary to introduce a pair of bulk spinors to describe baryons in QCD. The pair of spinors naturally interact with each other through a Yukawa coupling by a bulk scalar, corresponding to the order parameter of chiral symmetry breaking.

Our model has two adjustable parameters, the infrared cutoff z_m and the Yukawa coupling g , with which we fit the baryon spectrum. Taking the lowest-lying nucleon mass as an input, we have calculated the mass spectrum of excited nucleons and their couplings to pions, which are in a reasonable agreement with data, though it deviates from the data quite a lot for highly excited states. We have also calculated the pion-nucleon-nucleon couplings. The pion coupling with lowest-lying nucleons is about 50% less than the observed value, $g_{\pi NN} \simeq 13$. However we expect our results will improve if we take account of the anomalous dimension of the baryon operator and if we relax our AdS_5 geometry to have a soft IR wall.

We find that our model explains naturally the parity-doubling of excited baryons. The mass difference between parity-doublet nucleons and their couplings to pions become smaller for excited nucleons, suggesting that the parity-doubling or the effective chiral restoration in excited baryons is due to the holographic nature of QCD or large N_C QCD.

In our analysis we have focused on the spin- $\frac{1}{2}$ baryons in the two-flavor case. It is however straightforward to extend our work to spin- $\frac{3}{2}$ baryons and to the three-flavor case.

Acknowledgments

We acknowledge useful discussions with Nakwoo Kim, Youngman Kim, Shin Nakamura, Mannque Rho, Sang-Jin Sin and M.A.Stephanov. Two of the authors (D.K.H. and H.U.Y.) would like to thank the APCTP for organizing and sponsoring the focus program on “Search for Exotic State of Dense Matter” in 2006, where this work was partly done. The work of D. K. H. was supported by grant No. R01-2006-000-10912-0 from the Basic Research Program of the Korea Science & Engineering Foundation and H. U. Y. is partly supported by grant No. R01-2003-000-10391-0 from the Basic Research Program of the Korea Science & Engineering Foundation. The work of T.I. is supported by the grants of JSPS (Basic research B,C) and the Ministry of Education and Science (priority area).

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” Phys. Lett. B **428**, 105 (1998).
- [3] E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2**, 253 (1998).
- [4] J. Babington, J. Erdmenger, N. J. Evans, Z. Guralnik and I. Kirsch, “Chiral symmetry breaking and pions in non-supersymmetric gauge / gravity duals,” Phys. Rev. D **69**, 066007 (2004).
- [5] M. Kruczenski, D. Mateos, R. C. Myers and D. J. Winters, “Towards a holographic dual of large-N(c) QCD,” JHEP **0405**, 041 (2004).
- [6] T. Sakai and S. Sugimoto, “Low energy hadron physics in holographic QCD,” Prog. Theor. Phys. **113**, 843 (2005).
- [7] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, “QCD and a holographic model of hadrons,” Phys. Rev. Lett. **95**, 261602 (2005).
- [8] L. Da Rold and A. Pomarol, “Chiral symmetry breaking from five dimensional spaces,” Nucl. Phys. B **721**, 79 (2005).
- [9] N. Evans, J. P. Shock and T. Waterson, “Towards a perfect QCD gravity dual,” Phys. Lett. B **622**, 165 (2005).
- [10] K. Ghoroku, N. Maru, M. Tachibana and M. Yahiro, Phys. Lett. B **633**, 602 (2006).

- [11] D. K. Hong and H. U. Yee, “Holographic estimate of oblique corrections for technicolor,” Phys. Rev. D **74**, 015011 (2006).
- [12] J. Hirn and V. Sanz, “A negative S parameter from holographic technicolor,” [arXiv:hep-ph/0606086].
- [13] M. Piai, “Precision electro-weak parameters from AdS(5), localized kinetic terms and anomalous dimensions,” [arXiv:hep-ph/0608241].
- [14] C. Csaki, C. Grojean, L. Pilo and J. Terning, Phys. Rev. Lett. **92**, 101802 (2004).
- [15] G. Burdman and Y. Nomura, “Holographic theories of electroweak symmetry breaking without a Higgs boson,” Phys. Rev. D **69**, 115013 (2004).
- [16] G. F. de Teramond and S. J. Brodsky, “The hadronic spectrum of a holographic dual of QCD,” Phys. Rev. Lett. **94**, 201601 (2005).
- [17] M. Henningson and K. Sfetsos, “Spinors and the AdS/CFT correspondence,” Phys. Lett. B **431**, 63 (1998); W. Muck and K. S. Viswanathan, “Conformal field theory correlators from classical field theory on anti-de Sitter space. II: Vector and spinor fields,” Phys. Rev. D **58**, 106006 (1998); M. Henneaux, “Boundary terms in the AdS/CFT correspondence for spinor fields,” [arXiv:hep-th/9902137].
- [18] A. Volovich, “Rarita-Schwinger field in the AdS/CFT correspondence,” JHEP **9809**, 022 (1998); A. S. Koshelev and O. A. Rytchkov, “Note on the massive Rarita-Schwinger field in the AdS/CFT correspondence,” Phys. Lett. B **450**, 368 (1999); R. C. Rashkov, “Note on the boundary terms in AdS/CFT correspondence for Rarita-Schwinger field,” Mod. Phys. Lett. A **14**, 1783 (1999); P. Matlock and K. S. Viswanathan, “The AdS/CFT correspondence for the massive Rarita-Schwinger field,” Phys. Rev. D **61**, 026002 (2000).
- [19] R. L. Jaffe, D. Pirjol and A. Scardicchio, Phys. Rev. Lett. **96**, 121601 (2006); arXiv:hep-ph/0602010; L. Y. Glozman, arXiv:hep-ph/0609230.
- [20] G. ’t Hooft, “*Recent Developments In Gauge Theories*,” Proceedings, Nato Advanced Study (Plenum Press, New York 1980).
- [21] R. Contino and A. Pomarol, “Holography for fermions,” JHEP **0411**, 058 (2004).
- [22] R. Argurio, G. Ferretti and C. Petersson, “Massless fermionic bound states and the gauge / gravity correspondence,” JHEP **0603**, 043 (2006).
- [23] I. Kirsch, “Spectroscopy of fermionic operators in AdS/CFT,” [arXiv:hep-th/0607205].
- [24] D. T. Son and M. A. Stephanov, “QCD and dimensional deconstruction,” Phys. Rev. D **69**,

065020 (2004).

- [25] M. Eto, M. Nitta, K. Ohashi and D. Tong, “Skyrmions from instantons inside domain walls,”
Phys. Rev. Lett. **95**, 252003 (2005).